arXiv:1602.01496v1 [math.CA] 31 Jan 2016

SOME UNIFIED INTEGRALS ASSOCIATED WITH BESSEL-STRUVE KERNEL FUNCTION

K. S. NISAR, P. AGARWAL, AND S. JAIN

Abstract. In this paper, we discuss the generalized integral formula involving Bessel-Struve kernel function $S_\alpha(\lambda z)$, which expressed in terms of generalized Wright functions. Many interesting special cases also obtained in this study.

1. Introduction

In 1888 Pincherle studied the integrals involving product of Gamma functions along vertical lines (see [1, 2, 3]). Latterly, Barnes [4] , Mellin [5] and Cahen [6] extended the study and applied some of these integrals in the study of Riemann zeta function and other Drichlet's series. The integral formulas involving special functions have been developed by many researchers ([7],[8]). In [9] presented unified integral representation of Fox H-functions and in [10] hypergeometric ${}_2F_1$ functions. Recently J. Choi and P. Agarwal [11] obtained two unified integral representations of Bessel functions $J_v(z)$. Also, many interesting integral formula involving $J_v(z)$ is given in [7]and [12].

The Bessel-Struve kernel $S_\alpha(\lambda z), \lambda \in \mathbb{C}$, [14] which is unique solution of the initial value problem $l_\alpha u(z) = \lambda^2 u(z)$ with the initial conditions $u(0) = 1$ and $u'(0) = \lambda\Gamma(\alpha+1)/\sqrt{\pi}\Gamma(\alpha+3/2)$ is given by

$$S_\alpha(\lambda z) = j_\alpha(i\lambda z) - ih_\alpha(i\lambda z), \forall z \in C$$

where j_α and h_α are the normalized Bessel and Struve functions

Moreover, the Bessel-Struve kernel is a holomorphic function on $\mathbb{C}\times\mathbb{C}$ and it can be expanded in a power series in the form

$$S_\alpha(\lambda z) = \sum_{n=0}^{\infty} \frac{(\lambda z)^n \Gamma(\alpha+1)\Gamma((n+1)/2)}{\sqrt{\pi} n!\Gamma(n/2+\alpha+1)}, \tag{1.1}$$

The generalized Wright hypergeometric function ${}_p\psi_q(z)$ is given by the series

$${}_p\psi_q(z) = {}_p\psi_q\left[\begin{array}{c}(a_i,\alpha_i)_{1,p}\\(b_j,\beta_j)_{1,q}\end{array}\middle| z\right] = \sum_{k=0}^{\infty} \frac{\prod_{i=1}^{p}\Gamma(a_i+\alpha_i k)}{\prod_{j=1}^{q}\Gamma(b_j+\beta_j k)}\frac{z^k}{k!}, \tag{1.2}$$

where $a_i, b_j \in \mathbb{C}$, and real $\alpha_i, \beta_j \in \mathbb{R}$ $(i = 1, 2, \ldots, p; j = 1, 2, \ldots, q)$. Asymptotic behavior of this function for large values of argument of $z \in \mathbb{C}$ were studied in [21]

2000 *Mathematics Subject Classification.* Primary 05C38, 15A15; Secondary 05A15, 15A18.

Key words and phrases. Bessel Struve kernel function, generalized Wright functions,Integral representations.

and under the condition

$$\sum_{j=1}^{q}\beta_j-\sum_{i=1}^{p}\alpha_i>-1 \tag{1.3}$$

was found in the work of [22, 23]. Properties of this generalized Wright function were investigated in [25], (see also [26, 27]. In particular, it was proved [25] that ${}_p\psi_q(z)$, $z\in\mathbb{C}$ is an entire function under the condition (1.3).

The generalized hypergeometric function represented as follows [28]:

$${}_pF_q\left[\begin{array}{c}(\alpha_p);\\(\beta_q);\end{array}\ z\right]=\sum_{n=0}^{\infty}\frac{\Pi_{j=1}^{p}\left(\alpha_j\right)_n}{\Pi_{j=1}^{p}\left(\beta_j\right)_n}\frac{z^n}{n!}, \tag{1.4}$$

provided $p\leq q; p=q+1$ and $|z|<1$

where $(\lambda)_n$ is well known Pochhammer symbol defined for (for $\lambda\in C$) (see [28])

$$(\lambda)_n:=\begin{cases}1 & (n=0)\\ \lambda\left(\lambda+1\right)....\left(\lambda+n-1\right) & \left(n\in N:=\{1,2,3....\}\right)\end{cases} \tag{1.5}$$

$$(\lambda)_n=\frac{\Gamma\left(\lambda+n\right)}{\Gamma\left(\lambda\right)}\qquad\left(\lambda\in C\backslash Z_0^-\right). \tag{1.6}$$

where Z_0^- is the set of nonpositive integers.

If we put $\alpha_1=...=\alpha_p=\beta_1=....=\beta_q$ in (1.2),then (1.4) is a special case of the generalized Wright function:

$${}_p\psi_q(z)={}_p\psi_q\left[\begin{array}{c}(\alpha_1,1),...,(\alpha_p,1);\\(\beta_1,1),...,(\beta_q,1);\end{array}\ z\right]=\frac{\prod_{j=1}^{p}\Gamma(\alpha_j)}{\prod_{j=1}^{q}\Gamma(\beta_j)}\ {}_pF_q\left[\begin{array}{c}\alpha_1,...,\alpha_p;\\\beta_1,...,\beta_q;\end{array}\ z\right] \tag{1.7}$$

For the present investigation, we need the following result of Oberbettinger [30]

$$\int_0^{\infty}x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}dx=2\lambda a^{-\lambda}\left(\frac{a}{2}\right)^{\mu}\frac{\Gamma\left(2\mu\right)\Gamma\left(\lambda-\mu\right)}{\Gamma\left(1+\lambda+\mu\right)} \tag{1.8}$$

provided $0<Re\left(\mu\right)<Re\left(\lambda\right)$

Motivated by the work of [8] , here we present the integral formulas of Bessel-Struve Kernel function of first kind $S_\alpha\left(\lambda z\right)$,$\lambda\in\mathbb{C}$,which expressed interns of generalized Wright or generalized hypergeometric functions.

2. Main results

Two generalized integral formulas established here, which expressed in terms of generalized (Wright) hypergeometric functions (1.7) by inserting the Bessel-Struve kernel function of the first kind (1.1) with the suitable argument in the integrand of (1.8)

Theorem 1. *For $\lambda,\mu,\nu,\gamma\in\mathbb{C}$,and $x>0$, $\mathbb{R}\left(\lambda\right)>\mathbb{R}\left(\mu\right)>0$, then the following integral formula holds true:*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma y}{x+a+\sqrt{x^2+2ax}}\right)dx \\
&= \frac{2^{1-\mu}a^{\mu-\lambda}\Gamma(\alpha+1)\Gamma(2\mu)}{\sqrt{\pi}} \\
&\times {}_3\Psi_2\left[\begin{matrix}\left(\frac{1}{2},\frac{1}{2}\right),(\lambda+1,1),(\lambda-\mu,1); \\ (\lambda,1),(1+\lambda+\mu,1);\end{matrix}\ \frac{\gamma y}{a}\right] \qquad (2.1)
\end{aligned}$$

Proof. Consider the series representation of $S_\alpha\left(\frac{\gamma y}{x+a+\sqrt{x^2+2ax}}\right)$ and applying (1.8) . By interchanging the order of integration and summation,which verified by uniform convergence of the involved series under the given conditions, we get

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma y}{x+a+\sqrt{x^2+2ax}}\right)dx \\
&= \int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\sum_{n=0}^\infty \frac{\left(\frac{\gamma y}{x+a+\sqrt{x^2+2ax}}\right)^n \Gamma(\alpha+1)\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{n}{2}+\alpha+1\right)n!}dx \\
&= \sum_{n=0}^\infty \frac{(\gamma y)^n\Gamma(\alpha+1)\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{n}{2}+\alpha+1\right)n!}\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-(\lambda+n)}dx
\end{aligned}$$

in view of the conditions give in Theorem 1and applying the integral formula (1.8) ,we obtain the following integral representation:

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma y}{x+a+\sqrt{x^2+2ax}}\right)dx \\
&= 2^{1-\mu}a^{-\lambda+\mu}\Gamma(\alpha+1)\Gamma(2\mu)\pi^{-1/2}\sum_{n=0}^\infty \Gamma\left(\frac{n+1}{2}\right)\frac{\Gamma(\lambda+n+1)}{\Gamma(\lambda+n)} \\
&\times\frac{\Gamma(\lambda+n-\mu)}{\Gamma(\lambda+\mu+n+1)}\frac{\gamma^n y^n}{n!a^n}
\end{aligned}$$

which,upon using (1.7) ,yeilds (2.1). This completes proof of theorem 1 □

Theorem 2. *For $\lambda,\mu,\nu,\gamma\in\mathbb{C}$ with $0<\mathbb{R}(\mu)<\mathbb{R}(\lambda+\nu)$ and $x>0$. The following integral formula hold true:*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma xy}{x+a+\sqrt{x^2+2ax}}\right)dx \\
&= \frac{2^{1+\mu}a^{\mu-\lambda}\Gamma(\alpha+1)\Gamma(\lambda-\mu)}{\sqrt{\pi}\Gamma(1+\lambda+\mu)}\ {}_3\Psi_2\left[\begin{matrix}\left(\frac{1}{2},\frac{1}{2}\right),(2\mu,2),(\lambda+1,1); \\ (\lambda,1),\left(\alpha+1,\frac{1}{2}\right);\end{matrix}\right]\gamma y
\end{aligned}$$

Proof. Interchanging the order of integration and summation and the series representation of Bessel Struve kerenel function, we get

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma xy}{x+a+\sqrt{x^2+2ax}}\right)dx \\
= &\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} \\
&\times\sum_{n=0}^\infty\left(\frac{\gamma xy}{x+a+\sqrt{x^2+2ax}}\right)^n \frac{\Gamma(\alpha+1)\,\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{n}{2}+\alpha+1\right)n!}dx \\
= &\sum_{n=0}^\infty \frac{\gamma^n y^n \Gamma(\alpha+1)\,\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}\Gamma\left(\frac{n}{2}+\alpha+1\right)}\int_0^\infty x^{\mu+n-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-(\lambda+n)}dx
\end{aligned}$$

in view of the condition give in theorem 1,we can apply the integral formula (1.8) and obtain the following integral representation:

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} S_\alpha\left(\frac{\gamma xy}{x+a+\sqrt{x^2+2ax}}\right)dx \\
= &\frac{2^{1+\mu}a^{\mu-\lambda}\Gamma(\alpha+1)\,\Gamma(\lambda-\mu)}{\sqrt{\pi}\Gamma(1+\lambda+\mu)}\sum_{n=0}^\infty \Gamma\left(\frac{n+1}{2}\right)\frac{\Gamma(\lambda+n+1)\,\Gamma(2\mu+2n)}{\Gamma(\lambda+n)\,\Gamma\left(\frac{n}{2}+\alpha+1\right)}\frac{\gamma^n y^n}{n!a^n}
\end{aligned}$$

which gives the desired result. □

2.1. **Representation of Bessel Struve kernel function in terms of exponential function.** In this subsection we represent the Bessel Struve function in terms of exponential function. Also, we derive the Marichev Saigo Maeda operator representation of special cases. The representation Bessel Struve Kernel function interms of exponential function as:

$$S_{\frac{-1}{2}}(x) = e^x, \tag{2.2}$$

$$S_{\frac{1}{2}}(x) = \frac{-1+e^x}{x}. \tag{2.3}$$

Now, we give the the following corollaries:

Corollary 1. *For $\lambda,\mu\in\mathbb{C}$ with $0<\mathbb{R}(\mu)<\mathbb{R}(\lambda)$ and $x>0$. The following integral formula holds true*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} e^{\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)}dx \\
= &\; 2^{1-\mu}a^{\mu-\lambda}\Gamma(2\mu)\;{}_2\Psi_2\left[\begin{matrix}(\lambda+1,1),(\lambda-\mu,1);\\(\lambda,1),(1+\lambda-\mu);\end{matrix}\;\frac{y}{a}\right]
\end{aligned} \tag{2.4}$$

Proof. As same as in theorem 1 and theorem 2 , using the formula (1.8) and (2.2), one can easily reach the result □

Corollary 2. *Let the conditions given in Corollary 1 satisfied. Then the following integral formula holds true*

$$\begin{aligned}&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} e^{\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)}dx\\ =\quad &=\frac{2^{1-\mu}a^{\mu-\lambda}\Gamma(2\mu)\,\Gamma(\lambda+1)\,\Gamma(\lambda-\mu)}{\Gamma(\lambda)\,\Gamma(1+\lambda-\mu)}\;{}_2F_2\left[\begin{array}{l}\lambda+1,\lambda-\mu;\\ \lambda,1+\lambda-\mu;\end{array}\frac{y}{a}\right]\end{aligned}$$

Proof. In the view of equations (1.4) , (1.5) and (2.4) , we obtain the required result. □

Corollary 3. *For $\lambda,\mu,\in\mathbb{C}$ with $0<\mathbb{R}(\mu)<\mathbb{R}(\lambda)$ and $x>0$.The following integral formula holds true*

$$\begin{aligned}&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda} e^{\left(\frac{y}{x+a+\sqrt{x^2+2ax}}-1\right)}dx\\ =\quad &2^{-\mu}a^{\mu-\lambda}\Gamma(2\mu)\;{}_3\Psi_3\left[\begin{array}{l}\left(\frac{1}{2},\frac{1}{2}\right),(\lambda+1,1),(\lambda-\mu,1);\\ \left(\frac{1}{2},\frac{3}{2}\right),(\lambda,1),(1+\lambda+\mu,1);\end{array}\frac{y}{a}\right]\end{aligned}$$

2.2. **Relation between Bessel Struve kernel function and Bessel and Struve function of first kind.** In this subsection we show the relation between $S_\alpha(x)$ and Bessel function $I_v(x)$ and Struve function $L_v(x)$ by choosing particular values of α

$$S_0(x)=I_0(x)+L_0(x), \tag{2.5}$$

$$S_1(x)=\frac{2I_1(x)+L_1(x)}{x}, \tag{2.6}$$

In the light of above relations ,we have the following theorems:

Theorem 3. *For $\lambda,\mu\in\mathbb{C}$ with $0<\mathbb{R}(\mu)<\mathbb{R}(\lambda)$ and $x>0$.Then the following integral formula holds true:*

$$\begin{aligned}&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\\ &\times\left[I_0\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)+L_0\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)\right]dx\\ =\quad &2^{1-\mu}a^{\mu-\lambda}\pi^{-1/2}\Gamma(2\mu)\;{}_3\Psi_3\left[\begin{array}{l}\left(\frac{1}{2},\frac{1}{2}\right),(\lambda+1,1),(\lambda-\mu,1);\\ \left(1,\frac{1}{2}\right),(\lambda,1),(1+\lambda+\mu,1);\end{array}\frac{y}{a}\right]\end{aligned}$$

Proof. Consider the relation given in (2.5) and applying (1.8) . By interchanging the order of integration and summation,which verified by uniform convergence of

the involved series under the given conditions, we get

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\\
&\times\left[I_0\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)+L_0\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)\right]dx\\
=&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\sum_{n=0}^{\infty}\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)^{n}\frac{\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{\pi}n!\Gamma\left(\frac{n}{2}+1\right)}dx\\
=&\sum_{n=0}^{\infty}\frac{\Gamma\left(\frac{n+1}{2}\right)y^n}{\sqrt{\pi}n!\Gamma\left(\frac{n}{2}+1\right)}\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-(\lambda+n)}dx\\
=&\ 2^{1-\mu}a^{\mu-\lambda}\pi^{-1/2}\Gamma(2\mu)\sum_{n=0}^{\infty}\frac{\Gamma\left(\frac{n+1}{2}\right)y^n}{n!\Gamma\left(\frac{n}{2}+1\right)a^n}\frac{\Gamma(\lambda+n+1)}{\Gamma(\lambda+n)}\frac{\Gamma(\lambda+n-\mu)}{\Gamma(\lambda+n+1+\mu)}
\end{aligned}$$

which gives the desired result. □

Theorem 4. *The following integral formula holds true with* $\lambda,\mu,\in\mathbb{C}$ *with* $0<\mathbb{R}(\mu)<\mathbb{R}(\lambda)$ *and* $x>0$.

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\\
&\times\left[2I_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)+L_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)\right]dx\\
=&\ 2^{1-\mu}a^{\mu-\lambda}\pi^{-1/2}\Gamma(2\mu)\ {}_2\Psi_2\left[\begin{matrix}\left(\frac{1}{2},\frac{1}{2}\right),(\lambda-\mu,1);\\ \left(2,\frac{1}{2}\right),(1+\lambda+\mu,1);\end{matrix}\ \frac{y}{a}\right]
\end{aligned}$$

Proof. Consider the series representation of $S_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)$ and applying (1.8)and interchanging the order of integration and summation, we get

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\\
&\times\left[2I_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)+L_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)\right]dx\\
=&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}S_1\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)dx\\
=&\sum_{n=0}^{\infty}\frac{\Gamma\left(\frac{n+1}{2}\right)y^n}{\sqrt{\pi}n!\Gamma\left(\frac{n}{2}+2\right)}\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-(\lambda+n)}dx\\
=&\ 2^{1-\mu}a^{\mu-\lambda}\pi^{-1/2}\Gamma(2\mu)\sum_{n=0}^{\infty}\frac{\Gamma\left(\frac{n+1}{2}\right)\Gamma(\lambda+n-\mu)}{\Gamma\left(\frac{n}{2}+2\right)\Gamma(1+\lambda+n+\mu)}\frac{y^n}{a^n n!}
\end{aligned}$$

which gives the required result. □

Conclusion

The generalized integral formula involving Bessel-Struve kernel function $S_\alpha(\lambda z)$, which expressed in terms of generalized Wright functions are given in this paper. Also the relation between exponential function, Bessel function and Struve function with Bessel-Struve kernel function is also discussed with particular cases.

References

[1] S. Pincherle, Sulle funzioni ipergeometriche generalizzate Nota I, Atti della Reale Accademia dei Lincei. Rendiconti della Classe di Scienze Fisiche, Matematiche e Naturali, Serie 4, Vol. 4 (1888), 694–700

[2] S, Pincherle: Sulle funzioni ipergeometriche generalizzate. Nota I-II. In: Opere Scelte-I, Unione Matematica Italiana, Ed. Cremonese, Rome (1954), 223-239.

[3] Mainardi, F, Pagnini, G, Pincherle, S: The pioneer of the Mellin-Barnes integrals. J. Comput. Appl. Math. 153, (2003),331-342.

[4] Barnes, EW: A new development of the theory of the hypergeometric functions. Proc. Lond. Math. Soc. 6, 141-177, (1908)

[5] Mellin, HM: Abriß einer einheitlichen theorie der Gamma und der hypergeometrischen funktionen. Math. Ann. 68, 305-337 (1910)

[6] Cahen, E: Sur la fonction $\xi\sigma$ de Riemann et sur des fonctions analogues. Ann. Sci. Éc. Norm. Super. 11, 75-164, (1894)

[7] Brychkov, YA: Handbook of special functions: derivatives, integrals, series and other formulas. CRC Press, Boca Raton, (2008)

[8] Choi, J, Hasanov, A, Srivastava, HM, Turaev, M: Integral representations for Srivastava's triple hypergeometric functions. Taiwan. J. Math. 15, 2751-2762 (2011)

[9] Garg, M, Mittal, S: On a new unified integral. Proc. Indian Acad. Sci. Math. Sci. 114(2), 99-101 (2003)

[10] Ali, S: On some new unified integrals. Adv. Comput. Math. Appl. 1(3), 151-153 (2012)

[11] Junesang Choi, Praveen Agarwal, Certain unified integrals associated with Bessel functions, Boundary Value Problems 2013, 2013:95

[12] Watson, GN: A Treatise on the Theory of Bessel Functions. Cambridge Mathematical Library Edition. Camdridge University Press, Camdridge (1995). Reprinted (1996)

[13] H. Struve, Beitrag zur Theorie der Diffraction an Fernröhren, Ann. Physik Chemie, 17 (1882), 1008–1016.

[14] A. Gasmi and M. Sifi, The Bessel-Struve intertwinning operator on C and mean-periodic functions, IJMMS 2004:59, 3171–3185

[15] A. R. Ahmadi and S. E. Widnall, Unsteady lifting-line theory as a singular-perturbation problem. J. Fluid Mech. 153 (1985), 59-81.

[16] M. H. Hirata, Flow near the bow of a steadily turning ship, J. Fluid Mech. 71 (1975), no. 2, 283-291.

[17] D. C. Shaw, Perturbational results for di raction of water-waves by nearly-vertical barriers. IMA, J. Appl. Math. 34 (1985), no. 1, 99-117.

[18] J. Shao, P. Hanggi, Decoherent dynamics of a two-level system coupled to a sea of spins. Phys. Rev. Lett.81 (1998), no. 26, 5710-5713.

[19] J. Shao, P. Hanggi, Decoherent dynamics of a two-level system coupled to a sea of spins. Phys. Rev. Lett.81 (1998), no. 26, 5710-5713.

[20] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher transcendental functions. Vols. I, II*, McGraw-Hill Book Company, Inc., New York, 1953. NewYork, Toronto, London, 1953.

[21] C. Fox, The asymptotic expansion of generalized hypergeometric functions, Proc. London. Math. Soc. **27** (1928), no. 4, 389-400.

[22] E. M. Wright, The asymptotic expansion of integral functions defined by Taylor series, Philos. Trans. Roy. Soc. London, Ser. A. **238** (1940), 423–451.

[23] E. M. Wright, The asymptotic expansion of the generalized hypergeometric function, Proc. London Math. Soc. (2) **46** (1940), 389-408.

[24] A. A. Kilbas and M. Saigo,*H-transforms*, Chapman & Hall/CRC, Boca Raton, FL, 2004.

[25] A. A. Kilbas, M. Saigo and J. J. Trujillo, On the generalized Wright function, Fract. Calc. Appl. Anal. **5** (2002), no. 4, 437–460.

[26] A. A. Kilbas and N. Sebastian, Generalized fractional integration of Bessel function of the first kind, Integral Transforms Spec. Funct. **19** (2008), no. 11-12, 869–883.

[27] A. A. Kilbas and N. Sebastian, Fractional integration of the product of Bessel function of the first kind, Fract. Calc. Appl. Anal. **13** (2010), no. 2, 159–175.

[28] E. D. Rainville, *Special functions*, Macmillan, New York, 1960.

[29] Saiful. R. Mondal, Unified Integrals associated with generalized Bessel functions and Struve functions, Eng. Math. Let,10, 2015
[30] Oberhettinger, F: Tables of Mellin Transforms. Springer, New York (1974)

Department of Mathematics, College of Arts and Science-Wadi Addwasir, Prince Sattam bin Abdulaziz University, Saudi Arabia
E-mail address: ksnisar1@gmail.com

Department of Mathematics, Anand International College of Engineering, Jaipur 303012, Rajasthan, India.

Department of Mathematics, Poornima College of Engineering,Jaipur 302002, Rajasthan, India.